

# Tilapia ALLER PRIMO FLOAT



## Grower Feed



### DECLARATION

|                        | 3 mm | 4.5 mm | 6 mm |
|------------------------|------|--------|------|
| Crude protein (%)      | 37   | 37     | 37   |
| Crude fat (%)          | 12   | 12     | 12   |
| NFE (%)                | 35,2 | 35,2   | 35,2 |
| Ash (%)                | 6,2  | 6,2    | 6,2  |
| Fibre (%)              | 3,6  | 3,6    | 3,6  |
| P (%)                  | 1,0  | 1,0    | 1,0  |
| Gross energy (MJ)      | 20,1 | 20,1   | 20,1 |
| Digestible energy (MJ) | 15,4 | 15,4   | 15,4 |

### COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

blood products, fish meal, fish oil, grain products, marine by-products, processed animal proteins, vegetable oils, vegetable proteins, vitamins and minerals.

### RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

| Fish (g) | MM     | Water temperature (°C) |      |      |      |      |      |      |
|----------|--------|------------------------|------|------|------|------|------|------|
|          |        | 20                     | 22   | 24   | 26   | 28   | 30   | 32   |
| 70-100   | 3 mm   | 1,24                   | 1,86 | 2,32 | 2,79 | 3,1  | 2,79 | 2,48 |
| 100-200  | 3 mm   | 0,99                   | 1,49 | 1,86 | 2,23 | 2,48 | 2,23 | 1,98 |
| 200-400  | 4.5 mm | 0,79                   | 1,19 | 1,49 | 1,78 | 1,98 | 1,78 | 1,59 |
| 400-800  | 4.5 mm | 0,63                   | 0,95 | 1,19 | 1,43 | 1,59 | 1,43 | 1,27 |
| >800     | 6 mm   | 0,51                   | 0,76 | 0,95 | 1,14 | 1,27 | 1,14 | 1,01 |

### ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

|                  | 3 mm |      |      | 4.5 mm |      |      | 6 mm |      |      |
|------------------|------|------|------|--------|------|------|------|------|------|
| Feed conversion  | 1    | 1,1  | 1,2  | 1,1    | 1,2  | 1,3  | 1,2  | 1,3  | 1,4  |
| N in faeces (kg) | 0,47 | 0,52 | 0,57 | 0,52   | 0,57 | 0,62 | 0,57 | 0,62 | 0,66 |
| N in water (kg)  | 2,7  | 3,24 | 3,79 | 3,24   | 3,79 | 4,33 | 3,79 | 4,33 | 4,87 |
| P in faeces (kg) | 0,36 | 0,4  | 0,43 | 0,4    | 0,43 | 0,47 | 0,43 | 0,47 | 0,5  |
| P in water (kg)  | 0,41 | 0,49 | 0,58 | 0,49   | 0,58 | 0,66 | 0,58 | 0,66 | 0,75 |

### ENVIRONMENTAL IMPACT MEASURED IN CO<sub>2</sub>-EQ

Figures are in CO<sub>2</sub>-equivalents (kg/kg feed)

|                                              | 3 mm      | 4.5 mm    | 6 mm      |
|----------------------------------------------|-----------|-----------|-----------|
| CO <sub>2</sub> -eq. with land use change    | 1,06-1,48 | 1,06-1,48 | 1,06-1,48 |
| CO <sub>2</sub> -eq. without land use change | 0,88-1,05 | 0,88-1,05 | 0,88-1,05 |