

Tilapia - North Africa ALLER TIL-PRO OREA EX



Grower Feed



DECLARATION

| | 3 mm | 4.5 mm | 6 mm |
|------------------------|------|--------|------|
| Crude protein (%) | 33 | 33 | 33 |
| Crude fat (%) | 6 | 6 | 6 |
| NFE (%) | 44,8 | 44,8 | 44,8 |
| Ash (%) | 7,1 | 7,1 | 7,1 |
| Fibre (%) | 4,1 | 4,1 | 4,1 |
| P (%) | 1,2 | 1,2 | 1,2 |
| Gross energy (MJ) | 18,5 | 18,5 | 18,5 |
| Digestible energy (MJ) | 12,6 | 12,6 | 12,6 |

COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

DDGS, fish meal, fish oil, peas, rapeseed oil, single cell protein, soya, sunflower protein conc., triticale, vitamins and minerals, wheat.

RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

| Fish (g) | MM | Water temperature (°C) | | | | | | | | |
|----------|--------|------------------------|------|------|------|------|------|------|------|------|
| | | 16 | 18 | 20 | 22 | 24 | 26 | 28 | 30 | 32 |
| 70-100 | 3 mm | 0,58 | 0,97 | 1,55 | 2,32 | 2,9 | 3,48 | 3,87 | 3,48 | 3,1 |
| 100-200 | 3 mm | 0,46 | 0,77 | 1,24 | 1,86 | 2,32 | 2,79 | 3,1 | 2,79 | 2,48 |
| 200-400 | 4.5 mm | 0,37 | 0,62 | 0,99 | 1,49 | 1,86 | 2,23 | 2,48 | 2,23 | 1,98 |
| 400-800 | 4.5 mm | 0,3 | 0,5 | 0,79 | 1,19 | 1,49 | 1,78 | 1,98 | 1,78 | 1,58 |
| >800 | 6 mm | 0,24 | 0,4 | 0,63 | 0,95 | 1,19 | 1,43 | 1,58 | 1,43 | 1,27 |

ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

| | 3 mm | | | 4.5 mm | | | 6 mm | | |
|------------------|------|------|------|--------|------|------|------|------|------|
| Feed conversion | 1,2 | 1,3 | 1,4 | 1,3 | 1,4 | 1,5 | 1,4 | 1,5 | 1,6 |
| N in faeces (kg) | 0,51 | 0,55 | 0,59 | 0,55 | 0,59 | 0,63 | 0,59 | 0,63 | 0,68 |
| N in water (kg) | 3,08 | 3,56 | 4,05 | 3,56 | 4,05 | 4,54 | 4,05 | 4,54 | 5,02 |
| P in faeces (kg) | 0,44 | 0,48 | 0,52 | 0,48 | 0,52 | 0,56 | 0,52 | 0,56 | 0,59 |
| P in water (kg) | 0,61 | 0,69 | 0,78 | 0,69 | 0,78 | 0,87 | 0,78 | 0,87 | 0,95 |