

Tilapia - North Africa ALLER TIL-PRO 40% EX



Fry Feed



DECLARATION

	0.5 mm	1 mm	1.5 mm	2 mm
Crude protein (%)	40	40	40	40
Crude fat (%)	6	6	6	6
NFE (%)	33,8	33,8	33,8	33,8
Ash (%)	7,3	7,3	7,3	7,3
Fibre (%)	4,9	4,9	4,9	4,9
P (%)	0,9	0,9	0,9	0,9
Gross energy (MJ)	18,4	18,4	18,4	18,4
Digestible energy (MJ)	14,2	14,2	14,2	14,2

COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

corn gluten, d.l. methionine, DDGS, fish meal, fish oil, maize, monocalcium phosphate, soy oil, soya, sunflower meal, vitamins and minerals, wheat middlings, wheat products, yellow corn.

RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

Fish (g)	MM	Water temperature (°C)								
		16	18	20	22	24	26	28	30	32
0.5-1.0	0.5 mm	1,66	2,77	4,43	6,64	8,3	9,96	11,07	9,96	8,86
1.0-3.0	1 mm	1,36	2,27	3,63	5,44	6,8	8,17	9,07	8,17	7,26
3.0-6.0	1 mm	1,12	1,86	2,97	4,46	5,58	6,69	7,44	6,69	5,95
5.0-10.0	1.5 mm	0,91	1,52	2,44	3,66	4,57	5,49	6,1	5,49	4,88
10.0-30.0	2 mm	0,75	1,25	2	3	3,75	4,5	5	4,5	4
30.0-70.0	2 mm	0,61	1,02	1,64	1,46	3,07	3,69	4,1	3,69	3,28

ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

	0.5 mm			1 mm			1.5 mm			2 mm		
Feed conversion	0,9	1	1,1	1	1,1	1,2	1,1	1,2	1,3	1,2	1,3	1,4
N in faeces (kg)	0,46	0,51	0,56	0,51	0,56	0,61	0,56	0,61	0,67	0,61	0,67	0,72
N in water (kg)	2,55	3,14	3,73	3,14	3,73	4,32	3,73	4,32	4,9	4,32	4,9	5,49
P in faeces (kg)	0,23	0,26	0,28	0,26	0,28	0,31	0,28	0,31	0,33	0,31	0,33	0,36
P in water (kg)	0,19	0,25	0,31	0,25	0,31	0,37	0,31	0,37	0,43	0,37	0,43	0,49