

African catfish - North Africa ALLER CLARIA EX FLOAT, 2 MM



Pre-grower feed



DECLARATION

| | 2 mm |
|------------------------|------|
| Crude protein (%) | 45 |
| Crude fat (%) | 12 |
| NFE (%) | 25,0 |
| Ash (%) | 7,5 |
| Fibre (%) | 2,5 |
| P (%) | 1,1 |
| Gross energy (MJ) | 20,1 |
| Digestible energy (MJ) | 16,4 |

COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

corn gluten, DDGS, fish meal, fish oil, peas, rapeseed oil, single cell protein, soya, soya protein concentrate, sunflower protein conc., vitamins and minerals, wheat.

RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

| Fish (g) | MM | Water temperature (°C) | | | | | | | | |
|----------|------|------------------------|------|------|------|------|------|------|------|------|
| | | 14 | 16 | 18 | 20 | 22 | 24 | 26 | 28 | 30 |
| 10-30 | 2 mm | 0,85 | 1,42 | 2,27 | 3,4 | 4,25 | 5,1 | 5,67 | 5,1 | 4,54 |
| 30-50 | 2 mm | 0,72 | 1,21 | 1,93 | 2,89 | 3,62 | 4,34 | 4,82 | 4,34 | 3,86 |

ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

| | 2 mm | | |
|------------------|------|------|------|
| Feed conversion | 0,7 | 0,8 | 0,9 |
| N in faeces (kg) | 0,4 | 0,46 | 0,52 |
| N in water (kg) | 1,89 | 2,55 | 3,21 |
| P in faeces (kg) | 0,23 | 0,26 | 0,3 |
| P in water (kg) | 0,11 | 0,19 | 0,26 |