

# Sea bream - North Africa ALLER MARINE 48/15 EX



## Fry Feed



## DECLARATION

|                        | 1.5 mm | 2 mm |
|------------------------|--------|------|
| Crude protein (%)      | 48     | 48   |
| Crude fat (%)          | 15     | 15   |
| NFE (%)                | 19,6   | 19,6 |
| Ash (%)                | 7,1    | 7,1  |
| Fibre (%)              | 2,3    | 2,3  |
| P (%)                  | 1,1    | 1,1  |
| Gross energy (MJ)      | 21,0   | 21,0 |
| Digestible energy (MJ) | 16,0   | 16,0 |

## COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

corn gluten, DDGS, fish meal, fish oil, rice products, shrimpmeal, soy oil, sunflower meal, vitamins and minerals, wheat, wheat products.

## RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

| Fish (g) | MM     | Water temperature (°C) |      |      |      |      |      |      |      |      |
|----------|--------|------------------------|------|------|------|------|------|------|------|------|
|          |        | 12                     | 14   | 16   | 18   | 20   | 22   | 24   | 26   | 28   |
| 7-10     | 1.5 mm | 0,85                   | 1,2  | 1,54 | 1,88 | 2,22 | 2,73 | 3,25 | 3,42 | 2,91 |
| 10-15    | 2 mm   | 0,79                   | 1,1  | 1,49 | 1,97 | 2,63 | 3,51 | 4,16 | 4,38 | 3,94 |
| 15-20    | 2 mm   | 0,69                   | 0,96 | 1,31 | 1,74 | 2,31 | 3,09 | 3,66 | 3,86 | 3,47 |

## ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

|                  | 1.5 mm |      |      | 2 mm |      |      |
|------------------|--------|------|------|------|------|------|
| Feed conversion  | 0,8    | 0,9  | 1    | 0,9  | 1    | 1,1  |
| N in faeces (kg) | 0,49   | 0,55 | 0,61 | 0,55 | 0,61 | 0,68 |
| N in water (kg)  | 2,9    | 3,61 | 4,32 | 3,61 | 4,32 | 5,02 |
| P in faeces (kg) | 0,26   | 0,3  | 0,33 | 0,3  | 0,33 | 0,36 |
| P in water (kg)  | 0,19   | 0,26 | 0,34 | 0,26 | 0,34 | 0,42 |