

Carp ALLER LOOP



Grower Feed



DECLARATION

| | 3 mm |
|------------------------|------|
| Crude protein (%) | 40 |
| Crude fat (%) | 24 |
| NFE (%) | 20,1 |
| Ash (%) | 5,5 |
| Fibre (%) | 2,4 |
| P (%) | 1,1 |
| Gross energy (MJ) | 22,8 |
| Digestible energy (MJ) | 20,1 |

COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

blood products, fish meal, fish oil, grain products, marine by-products, non-marine by-products, processed animal proteins, single cell proteins, vegetable oils, vegetable proteins, vitamins and minerals.

RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

| Fish (g) | MM | Water temperature (°C) | | | | |
|----------|------|------------------------|------|------|------|------|
| | | 20 | 22 | 24 | 26 | 28 |
| 50-100 | 3 mm | 3,1 | 3,72 | 4,13 | 3,72 | 3,31 |
| 100-300 | 3 mm | 2,48 | 2,98 | 3,31 | 2,98 | 2,65 |

ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

| | 3 mm | | |
|------------------|------|------|------|
| Feed conversion | 0,9 | 1 | 1,1 |
| N in faeces (kg) | 0,46 | 0,51 | 0,56 |
| N in water (kg) | 2,55 | 3,14 | 3,73 |
| P in faeces (kg) | 0,3 | 0,33 | 0,36 |
| P in water (kg) | 0,26 | 0,34 | 0,42 |

ENVIRONMENTAL IMPACT MEASURED IN CO₂-EQ

Figures are in CO₂-equivalents (kg/kg feed)

| | 3 mm |
|--|-----------|
| CO ₂ -eq. with land use change | 1,34-1,77 |
| CO ₂ -eq. without land use change | 1,09-1,19 |