

Tilapia - West Africa
ALLER TIL-PRO OREA

Grower Feed

DECLARATION

	3 mm	4.5 mm	6 mm
Crude protein (%)	33	33	33
Crude fat (%)	6	6	6
NFE (%)	47,9	47,9	47,9
Ash (%)	5,2	5,2	5,2
Fibre (%)	2,9	2,9	2,9
P (%)	1	1	1
Gross energy (MJ)	18,9	18,9	18,9
Digestible energy (MJ)	15,0	15,0	15,0

COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

blood products, fish meal, fish oil, grain products, marine by-products, processed animal oils, processed animal proteins, vegetable oils, vegetable proteins, vitamins and minerals.

RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

		Water temperature (°C)					
Fish (g)	MM	22	24	26	28	30	32
70-100	3 mm	1,95	2,43	2,92	3,24	2,92	2,59
100-200	3 mm	1,56	1,95	2,33	2,59	2,33	2,08
200-400	4.5 mm	1,25	1,56	1,87	2,08	1,87	1,66
400-800	4.5 mm	1	1,25	1,49	1,66	1,49	1,33
>800	6 mm	0,8	1	1,2	1,33	1,2	1,06

ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

	3 mm			4.5 mm			6 mm		
Feed conversion	1,2	1,3	1,4	1,3	1,4	1,5	1,4	1,5	1,6
N in faeces (kg)	0,51	0,55	0,59	0,55	0,59	0,63	0,59	0,63	0,68
N in water (kg)	3,08	3,56	4,05	3,56	4,05	4,54	4,05	4,54	5,02
P in faeces (kg)	0,36	0,39	0,42	0,39	0,42	0,45	0,42	0,45	0,48
P in water (kg)	0,41	0,48	0,55	0,48	0,55	0,62	0,55	0,62	0,69

ENVIRONMENTAL IMPACT MEASURED IN CO_2-EQ

Figures are in CO_2-equivalents (kg/kg feed)

	3 mm	4.5 mm	6 mm
CO_2-eq. with land use change	0,97-1,58	0,97-1,58	0,97-1,58
CO_2-eq. without land use change	0,78-0,89	0,78-0,89	0,78-0,89

19/03/2024