

Sturgeon ALLER BRONZE EX



Grower Feed



DECLARATION

| | 3 mm | 4.5 mm | 6 mm | 8 mm |
|------------------------|------|--------|------|------|
| Crude protein (%) | 45 | 45 | 45 | 45 |
| Crude fat (%) | 15 | 15 | 15 | 15 |
| NFE (%) | 24,0 | 24,0 | 24,0 | 24,0 |
| Ash (%) | 6,9 | 6,9 | 6,9 | 6,9 |
| Fibre (%) | 3,1 | 3,1 | 3,1 | 3,1 |
| P (%) | 1,0 | 1,0 | 1,0 | 1,0 |
| Gross energy (MJ) | 21,2 | 21,2 | 21,2 | 21,2 |
| Digestible energy (MJ) | 17,2 | 17,2 | 17,2 | 17,2 |

COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

fish meal, fish oil, grain products, marine by-products, non-marine by-products, single cell proteins, vegetable oils, vegetable proteins, vitamins and minerals.

RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

| Fish (g) | MM | Water temperature (°C) | | | | | | | | |
|-----------|--------|------------------------|------|------|------|------|------|------|------|------|
| | | 10 | 12 | 14 | 16 | 18 | 20 | 22 | 24 | 26 |
| 50-100 | 3 mm | 0,2 | 0,71 | 1,22 | 1,63 | 2,04 | 1,83 | 1,63 | 1,02 | 0,2 |
| 100-200 | 3 mm | 0,17 | 0,59 | 1,01 | 1,35 | 1,69 | 1,52 | 1,35 | 0,84 | 0,17 |
| 200-800 | 4.5 mm | 0,14 | 0,49 | 0,84 | 1,12 | 1,4 | 1,26 | 1,12 | 0,7 | 0,14 |
| 800-1500 | 4.5 mm | 0,12 | 0,41 | 0,7 | 0,93 | 1,16 | 1,05 | 0,93 | 0,58 | 0,12 |
| 1500-3000 | 6 mm | 0,1 | 0,34 | 0,58 | 0,77 | 0,97 | 0,87 | 0,77 | 0,48 | 0,1 |
| 3000-4000 | 6 mm | 0,08 | 0,28 | 0,48 | 0,64 | 0,8 | 0,72 | 0,64 | 0,4 | 0,08 |
| 4000-7000 | 8 mm | 0,07 | 0,23 | 0,4 | 0,53 | 0,67 | 0,6 | 0,53 | 0,33 | 0,07 |

ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

| | 3 mm | | | 4.5 mm | | | 6 mm | | | 8 mm | | |
|------------------|------|------|------|--------|------|------|------|------|------|------|------|------|
| Feed conversion | 1 | 1,1 | 1,2 | 1,1 | 1,2 | 1,3 | 1,2 | 1,3 | 1,4 | 1,3 | 1,4 | 1,5 |
| N in faeces (kg) | 0,58 | 0,63 | 0,69 | 0,63 | 0,69 | 0,75 | 0,69 | 0,75 | 0,81 | 0,75 | 0,81 | 0,86 |
| N in water (kg) | 3,87 | 4,54 | 5,2 | 4,54 | 5,2 | 5,86 | 5,2 | 5,86 | 6,52 | 5,86 | 6,52 | 7,19 |
| P in faeces (kg) | 0,3 | 0,33 | 0,36 | 0,33 | 0,36 | 0,39 | 0,36 | 0,39 | 0,42 | 0,39 | 0,42 | 0,45 |
| P in water (kg) | 0,27 | 0,34 | 0,41 | 0,34 | 0,41 | 0,48 | 0,41 | 0,48 | 0,55 | 0,48 | 0,55 | 0,62 |

ENVIRONMENTAL IMPACT MEASURED IN CO₂-EQ

Figures are in CO₂-equivalents (kg/kg feed)

| | 3 mm | 4.5 mm | 6 mm | 8 mm |
|--|-----------|-----------|-----------|-----------|
| CO ₂ -eq. with land use change | 1,47-1,94 | 1,47-1,94 | 1,47-1,94 | 1,47-1,94 |
| CO ₂ -eq. without land use change | 1,08-1,21 | 1,08-1,21 | 1,08-1,21 | 1,08-1,21 |

18/05/2024