

Tilapia - East Africa ALLER TIL-PRO CAGE



Grower Feed



DECLARATION

| | 3 mm | 4 mm | 5 mm |
|------------------------|------|------|------|
| Crude protein (%) | 36 | 34 | 34 |
| Crude fat (%) | 5 | 4 | 4 |
| NFE (%) | 39,1 | 41,7 | 41,7 |
| Ash (%) | 6,9 | 7,3 | 7,3 |
| Fibre (%) | 5 | 5 | 5 |
| P (%) | 1,2 | 1,2 | 1,2 |
| Gross energy (MJ) | 18 | 17,6 | 17,6 |
| Digestible energy (MJ) | 16,1 | 15,7 | 15,7 |

COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

blood products, corn gluten, fish meal, fish oil, maize, meat & bone meal, soy oil, soya, vitamins and minerals, wheat products.

RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

| Fish (g) | MM | Water temperature (°C) | | | | | | | | |
|----------|------|------------------------|------|------|------|------|------|------|------|------|
| | | 16 | 18 | 20 | 22 | 24 | 26 | 28 | 30 | 32 |
| 70-100 | 3 mm | 0,46 | 0,77 | 1,24 | 1,86 | 2,32 | 2,79 | 3,1 | 2,79 | 2,48 |
| 100-200 | 3 mm | 0,37 | 0,62 | 0,99 | 1,49 | 1,86 | 2,23 | 2,48 | 2,23 | 1,98 |
| 200-400 | 4 mm | 0,3 | 0,5 | 0,79 | 1,19 | 1,49 | 1,78 | 1,98 | 1,78 | 1,59 |
| 400-600 | 4 mm | 0,24 | 0,41 | 0,65 | 0,97 | 1,22 | 1,46 | 1,62 | 1,46 | 1,3 |
| >600 | 5 mm | 0,19 | 0,32 | 0,52 | 0,78 | 0,97 | 1,17 | 1,3 | 1,17 | 1,04 |

ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

| | 3 mm | 4 mm | 5 mm |
|------------------|------|------|------|
| Feed conversion | 1,3 | 1,4 | 1,5 |
| N in faeces (kg) | 0,6 | 0,65 | 0,69 |
| N in water (kg) | 4,14 | 4,67 | 5,2 |
| P in faeces (kg) | 0,47 | 0,51 | 0,54 |
| P in water (kg) | 0,75 | 0,83 | 0,91 |