

# Perch

## ALLER INFA EX GR, 0.1-0.4 MM



### Fry Feed



#### DECLARATION

	0.1 mm	0.2 mm	0.4 mm
Crude protein (%)	64	64	64
Crude fat (%)	12	12	12
NFE (%)	4,9	4,9	4,9
Ash (%)	12,1	12,1	12,1
Fibre (%)	1,0	1,0	1,0
P (%)	1,4	1,4	1,4
Gross energy (MJ)	20,8	20,8	20,8
Digestible energy (MJ)	19,6	19,6	19,6

#### COMPOSITION

Raw materials listed alphabetically. The full composition will appear on the label

fish meal, fish oil, functional ingredients, grain products, krill meal, vegetable proteins, vitamins and minerals.

#### RECOMMENDED FEEDING LEVELS

Kg feed per 100 kg fish per day

Fish (g)	Granulate	Water temperature (°C)									
			14	16	18	20	22	24	26	28	30
0.05-0.2	0.1	0.1	1,18	1,97	3,15	4,73	5,91	7,1	7,89	7,1	6,31
0.2-0.5	0.2	0.2	0,99	1,64	2,63	3,94	4,93	5,91	6,57	5,91	5,26
0.5-1.0	0.4	0.4	0,82	1,37	2,19	3,29	4,11	4,93	5,48	4,93	4,38

#### ENVIRONMENTAL IMPACT WITH EXEMPLARY FEED CONVERSION RATIOS

Figures are per 100 kg fish production

	0.1 mm			0.2 mm			0.4 mm		
Feed conversion	0,5	0,6	0,7	0,6	0,7	0,8	0,7	0,8	0,9
N in faeces (kg)	0,31	0,37	0,43	0,37	0,43	0,49	0,43	0,49	0,55
N in water (kg)	2,06	3,03	3,99	3,03	3,99	4,95	3,99	4,95	5,91
P in faeces (kg)	0,23	0,27	0,32	0,27	0,32	0,36	0,32	0,36	0,41
P in water (kg)	0,1	0,2	0,31	0,2	0,31	0,41	0,31	0,41	0,52

#### ENVIRONMENTAL IMPACT MEASURED IN CO<sub>2</sub>-EQ

Figures are in CO<sub>2</sub>-equivalents (kg/kg feed)

	0.1 mm	0.2 mm	0.4 mm
CO <sub>2</sub> -eq. with land use change	1,14-1,73	1,14-1,73	1,14-1,73
CO <sub>2</sub> -eq. without land use change	1,10-1,67	1,10-1,67	1,10-1,67